

# Wavelet analysis as a $p$ -adic spectral analysis

S.V.Kozyrev

February 3, 2008

*Institute of Chemical Physics, Russian Academy of Science*

## Abstract

New orthonormal basis of eigenfunctions for the Vladimirov operator of  $p$ -adic fractional derivation is constructed. The map of  $p$ -adic numbers onto real numbers ( $p$ -adic change of variable) is considered.  $p$ -Adic change of variable maps the Haar measure on  $p$ -adic numbers onto the Lebesgue measure on the positive semiline.  $p$ -Adic change of variable (for  $p = 2$ ) provides an equivalence between the constructed basis of eigenfunctions of the Vladimirov operator and the wavelet basis in  $L^2(\mathbf{R}_+)$  generated from the Haar wavelet. This means that the wavelet analysis can be considered as a  $p$ -adic spectral analysis.

## 1 Introduction

In the present paper we construct a new orthonormal basis of eigenfunctions of the Vladimirov operator of  $p$ -adic fractional derivation. The example of such a basis one can find in [1]. Different bases of eigenvectors of the Vladimirov operator were built in [2], [3], [4]. The basis constructed in the present paper consists of locally constant functions with support on  $p$ -adic discs.

We also check that the constructed basis of eigenfunctions of the Vladimirov operator (for  $p = 2$ ) is equivalent to the wavelet basis in  $L^2(\mathbf{R}_+)$  generated from the Haar wavelet. This equivalence is given by  $p$ -adic change of variables: the map of  $p$ -adic numbers onto real numbers that conserves the measure. This means that the wavelet analysis can be considered as a  $p$ -adic

harmonic analysis (decomposition of functions over the eigenfunctions of the Vladimirov operator of  $p$ -adic fractional derivation).

For introduction to  $p$ -adic analysis see [1].  $p$ -Adic analysis and  $p$ -adic mathematical physics attract great interest, see [1], [5], [6]. For instance,  $p$ -adic models in string theory were introduced, see [7], [8], and  $p$ -adic quantum mechanics [9] and  $p$ -adic quantum gravity [10] were investigated.  $p$ -Adic analysis was applied to investigate the spontaneous breaking of the replica symmetry, cf. [11], [12], [13], [14].

Let us make here a brief review of  $p$ -adic analysis. The field  $Q_p$  of  $p$ -adic numbers is the completion of the field of rational numbers  $Q$  with respect to the  $p$ -adic norm on  $Q$ . This norm is defined in the following way. An arbitrary rational number  $x$  can be written in the form  $x = p^\gamma \frac{m}{n}$  with  $m$  and  $n$  not divisible by  $p$ . The  $p$ -adic norm of the rational number  $x = p^\gamma \frac{m}{n}$  is equal to  $|x|_p = p^{-\gamma}$ .

The most interesting property of the field of  $p$ -adic numbers is ultrametricity. This means that  $Q_p$  obeys the strong triangle inequality

$$|x + y|_p \leq \max(|x|_p, |y|_p).$$

We will consider disks in  $Q_p$  of the form  $\{x \in Q_p : |x - x_0|_p \leq p^{-k}\}$ . For example, the ring  $Z_p$  of integer  $p$ -adic numbers is the disk  $\{x \in Q_p : |x|_p \leq 1\}$  which is the completion of integers with the  $p$ -adic norm. The main properties of disks in arbitrary ultrametric space are the following:

1. Every point of a disk is the center of this disk.
2. Two disks either do not intersect or one of these disks contains the other.

The  $p$ -adic Fourier transform  $F$  of the function  $f(x)$  is defined as follows

$$F[f](\xi) = \tilde{f}(\xi) = \int_{Q_p} \chi(\xi x) f(x) d\mu(x)$$

Here  $d\mu(x)$  is the Haar measure. The inverse Fourier transform have the form

$$F^{-1}[\tilde{g}](x) = \int_{Q_p} \chi(-\xi x) \tilde{g}(\xi) d\mu(\xi)$$

Here  $\chi(\xi x) = \exp(2\pi i \xi x)$  is the character of the field of  $p$ -adic numbers. For example, the Fourier transform of the indicator function  $\Omega(x)$  of the disk of

radius 1 with center in zero (this is a function that equals to 1 on the disk and to 0 outside the disk) is the function of the same type:

$$\tilde{\Omega}(\xi) = \Omega(\xi)$$

In the present paper we use the following Vladimirov operator  $D_x^\alpha$  of the fractional  $p$ -adic differentiation, that is defined [1] as

$$D_x^\alpha f(x) = F^{-1} \circ |\xi|_p^\alpha \circ F[f](x) = \frac{p^\alpha - 1}{1 - p^{-1-\alpha}} \int_{Q_p} \frac{f(x) - f(y)}{|x - y|_p^{1+\alpha}} d\mu(y) \quad (1)$$

Here  $F$  is the ( $p$ -adic) Fourier transform, the second equality holds for  $\alpha > 0$ .

For further reading on the subject of  $p$ -adic analysis see [1].

## 2 $p$ -Adic spectral analysis

Let us construct an orthonormal basis of eigenvectors of the Vladimirov operator. The following lemma gives the basic technical result.

**Lemma 1.** *The function*

$$\psi(x) = \chi(p^{-1}x)\Omega(|x|_p) \quad (2)$$

*is an eigenvector of the Vladimirov operator:*

$$D^\alpha \psi(x) = p^\alpha \psi(x)$$

**Remark.** The eigenvalue for  $\psi(x)$  is the same as for the character  $\chi(p^{-1}x)$ :

$$D^\alpha \chi(p^{-1}x) = p^\alpha \chi(p^{-1}x)$$

*Proof* Let us check that  $\psi(x)$  is an eigenvector of the Vladimirov operator. We get

$$D^\alpha \psi(x) = \frac{p^\alpha - 1}{1 - p^{-1-\alpha}} \int \frac{\psi(x) - \psi(y)}{|x - y|_p^{1+\alpha}} d\mu(y) =$$

$$\begin{aligned}
&= \frac{p^\alpha - 1}{1 - p^{-1-\alpha}} \int \frac{\chi(p^{-1}x)\Omega(|x|_p) - \chi(p^{-1}y)\Omega(|y|_p)}{|x - y|_p^{1+\alpha}} d\mu(y) = \\
&= \chi(p^{-1}x) \frac{p^\alpha - 1}{1 - p^{-1-\alpha}} \int \frac{\Omega(|x|_p) - \chi(p^{-1}(y - x))\Omega(|y|_p)}{|x - y|_p^{1+\alpha}} d\mu(y) \quad (3)
\end{aligned}$$

Let us calculate the integral over  $y$  in (3) in two cases.

1) Let  $|x|_p \leq 1$ . In this case the integral in (3) is given by

$$\int \frac{1 - \chi(p^{-1}(y - x))\Omega(|y|_p)}{|x - y|_p^{1+\alpha}} d\mu(y)$$

Using that every point of  $p$ -adic disk is its center we get that for  $|x|_p \leq 1$  we have  $\Omega(|y|_p) = \Omega(|x - y|_p)$  and therefore the integral is equal to

$$\int \frac{1 - \chi(p^{-1}(y - x))\Omega(|y - x|_p)}{|x - y|_p^{1+\alpha}} d\mu(y) = \int \frac{1 - \chi(p^{-1}z)\Omega(|z|_p)}{|z|_p^{1+\alpha}} d\mu(z)$$

2) Let  $|x|_p > 1$ . For the integral in (3) we get

$$-\frac{1}{|x|_p^{1+\alpha}} \int_{|y|_p \leq 1} \chi(p^{-1}(y - x)) d\mu(y) = 0$$

This proves that  $\psi(x)$  is an eigenvector of the Vladimirov operator with the following eigenvalue

$$\begin{aligned}
D^\alpha \psi(x) &= \psi(x) \frac{p^\alpha - 1}{1 - p^{-1-\alpha}} \int \frac{1 - \chi(p^{-1}z)\Omega(|z|_p)}{|z|_p^{1+\alpha}} d\mu(z) = \\
&= \psi(x) \frac{p^\alpha - 1}{1 - p^{-1-\alpha}} \left( p^{-1} \sum_{i=0}^{p-1} (1 - \chi(p^{-1}i)) + (1 - p^{-1}) \sum_{\gamma=1}^{\infty} p^\gamma p^{-(1+\alpha)\gamma} \right) = p^\alpha \psi(x)
\end{aligned}$$

that finishes the proof of the lemma.

The lemma implies

$$D^\alpha \psi(ax + b) = |a|_p^\alpha p^\alpha \psi(ax + b) \quad (4)$$

One can check that the set of (integrable) functions  $\psi(ax + b)$  (for different  $a, b$ ) is a complete system in a Hilbert space  $L^2(Q_p)$ . Moreover,

**Theorem 2.** The set of functions  $\{\psi_{\gamma jn}\}$ :

$$\psi_{\gamma jn}(x) = p^{-\frac{\gamma}{2}} \chi(p^{\gamma-1} jx) \Omega(|p^\gamma x - n|_p), \quad \gamma \in \mathbf{Z}, \quad n \in Q_p/Z_p, \quad j = 1, \dots, p-1 \quad (5)$$

is an orthonormal basis in  $L^2(Q_p)$  of eigenvectors of the operator  $D^\alpha$ :

$$D^\alpha \psi_{\gamma jn} = p^{\alpha(1-\gamma)} \psi_{\gamma jn} \quad (6)$$

The group  $Q_p/Z_p$  in (5) is parameterized by

$$n = \sum_{i=1}^m n_i p^{-i}, \quad n_i = 0, \dots, p-1$$

*Proof* Consider the scalar product

$$\begin{aligned} & \langle \psi_{\gamma jn}, \psi_{\gamma' j'n'} \rangle = \\ & = \int_{Q_p} p^{-\frac{\gamma}{2}} \chi(-p^{\gamma-1} jx) \Omega(|p^\gamma x - n|_p) p^{-\frac{\gamma'}{2}} \chi(p^{\gamma'-1} j'x) \Omega(|p^{\gamma'} x - n'|_p) d\mu(x) \quad (7) \end{aligned}$$

Consider  $\gamma \leq \gamma'$ . We have that the product of indicators is equal to the indicator or zero:

$$\Omega(|p^\gamma x - n|_p) \Omega(|p^{\gamma'} x - n'|_p) = \Omega(|p^\gamma x - n|_p) \Omega(|p^{\gamma'-\gamma} n - n'|_p)$$

Since  $n' \in Q_p/Z_p$  the function  $\Omega(|p^{\gamma'-\gamma} n - n'|_p)$  is non-zero (and equal to one) for

$$n' = p^{\gamma'-\gamma} n$$

We get for (7)

$$\int_{Q_p} p^{-\frac{\gamma}{2}} \chi(-p^{\gamma-1} jx) \Omega(|p^\gamma x - n|_p) p^{-\frac{\gamma'}{2}} \chi(p^{\gamma'-1} j'x) \Omega(|p^\gamma x - n|_p) \Omega(|p^{\gamma'-\gamma} n - n'|_p) d\mu(x) \quad (8)$$

Consider  $\gamma < \gamma'$ . Then for the integral (8) we get

$$\int_{Q_p} p^{-\frac{\gamma}{2}} p^{-\frac{\gamma'}{2}} \chi(-p^{\gamma-1} jx) \chi(p^{\gamma'-\gamma-1} j'n) \Omega(|p^\gamma x - n|_p) \Omega(|p^{\gamma'-\gamma} n - n'|_p) d\mu(x) = 0$$

Therefore the scalar product (7) can be non-zero only for  $\gamma = \gamma'$ . For  $\gamma = \gamma'$  the integral (8) is equal to

$$\int_{Q_p} p^{-\gamma} \chi(-p^{\gamma-1} j x) \chi(p^{\gamma-1} j' x) \Omega(|p^\gamma x - n|_p) \Omega(|n - n'|_p) d\mu(x)$$

Since  $n, n' \in Q_p/Z_p$

$$\Omega(|n - n'|_p) = \delta_{nn'}$$

we get for (7)

$$\begin{aligned} \langle \psi_{\gamma j n}, \psi_{\gamma' j' n'} \rangle &= \delta_{\gamma\gamma'} \delta_{nn'} \int_{Q_p} p^{-\gamma} \chi(p^{\gamma-1} (j' - j) x) \Omega(|p^\gamma x - n|_p) d\mu(x) = \\ &= \delta_{\gamma\gamma'} \delta_{nn'} \delta_{jj'} \end{aligned}$$

that proves that the vectors  $\psi_{\gamma j n}$  are orthonormal.

To prove that the set of vectors  $\{\psi_{\gamma j n}\}$  is an orthonormal basis (is total in  $L^2(Q_p)$ ) we use the Parsevale identity.

Since the set of indicators (characteristic functions) of  $p$ -adic discs is total in  $L^2(Q_p)$  and the set of vectors  $\{\psi_{\gamma j n}\}$  is translationally invariant and invariant under dilations  $x \mapsto p^n x$ ,  $x \in Q_p$ , to prove that  $\{\psi_{\gamma j n}\}$  is a complete system it is enough to check the Parsevale identity for the indicator  $\Omega(|x|_p)$ .

We have

$$\langle \Omega(|x|_p), \psi_{\gamma j n} \rangle = p^{-\frac{\gamma}{2}} \theta(\gamma) \delta_{n0}, \quad \theta(\gamma) = 0, \gamma \leq 0, \quad \theta(\gamma) = 1, \gamma \geq 1 \quad (9)$$

Formula (9) implies the Parsevale identity for  $\Omega(|x|_p)$ :

$$\sum_{\gamma j n} |\langle \Omega(|x|_p), \psi_{\gamma j n} \rangle|^2 = \sum_{\gamma=1}^{\infty} (p-1) p^{-\gamma} = 1 = |\langle \Omega(|x|_p), \Omega(|x|_p) \rangle|^2$$

that proves that  $\{\psi_{\gamma j n}\}$  is an orthonormal basis in  $L^2(Q_p)$ .

Formula (4) implies that the basis  $\{\psi_{\gamma j n}\}$  is an orthonormal basis of eigenvectors of the operator  $D^\alpha$  with eigenvalues (6). This finishes the proof of the theorem.

### 3 Wavelet interpretation

Let us discuss the connection between the constructed basis  $\{\psi_{\gamma j n}\}$  and the basis of wavelets in the space of quadratically integrable functions  $L^2(\mathbf{R}_+)$  on positive semiline. The wavelet basis in  $L^2(\mathbf{R}_+)$  is a basis given by shifts and dilations of the so called mother wavelet function, cf. [15]. The simplest example of such a function is the Haar wavelet

$$\Psi(x) = \chi_{[0, \frac{1}{2}]}(x) - \chi_{[\frac{1}{2}, 1]}(x) \quad (10)$$

(difference of two characteristic functions).

The wavelet basis in  $L^2(\mathbf{R})$  (or basis of multiresolution wavelets) is the basis

$$\Psi_{\gamma n}(x) = 2^{-\frac{\gamma}{2}} \Psi(2^{-\gamma} x - n), \quad \gamma \in \mathbf{Z}, \quad n \in \mathbf{Z} \quad (11)$$

Consider the  $p$ -adic change of variables, i.e. the onto map

$$\begin{aligned} \rho : Q_p &\rightarrow \mathbf{R}_+ \\ \rho : \sum_{i=\gamma}^{\infty} a_i p^i &\mapsto \sum_{i=\gamma}^{\infty} a_i p^{-i-1}, \quad a_i = 0, \dots, p-1, \quad \gamma \in \mathbf{Z} \end{aligned} \quad (12)$$

This map is not a one-to-one map. The map  $\rho$  is continuous and moreover

**Lemma 3.** *The map  $\rho$  satisfies the Holder inequality*

$$|\rho(x) - \rho(y)| \leq |x - y|_p \quad (13)$$

*Proof* Consider

$$x = \sum_{i=\alpha}^{\infty} x_i p^i, \quad y = \sum_{i=\beta}^{\infty} y_i p^i,$$

where  $\alpha \leq \beta$ . Then

$$\rho(x) - \rho(y) = \sum_{i=\alpha}^{\beta-1} x_i p^i + \sum_{i=\beta}^{\infty} (x_i - y_i) p^i$$

Consider the following two cases:

1) Let  $\alpha < \beta$ . Then

$$|\rho(x) - \rho(y)| \leq (p-1) \sum_{i=\alpha}^{\infty} p^{-i-1} = |x - y|_p$$

2) Let  $\alpha = \beta$ . Then  $|x - y|_p = p^{-\gamma}$ ,  $\gamma > \alpha$ .

$$\rho(x) - \rho(y) = \sum_{i=\gamma}^{\infty} (x_i - y_i) p^{-i-1}$$

$$|\rho(x) - \rho(y)| \leq (p-1) \sum_{i=\gamma}^{\infty} p^{-i-1} = p^{-\gamma} = |x - y|_p$$

that finishes the proof of the lemma.

The following map is a one-to-one map:

$$\rho : Q_p/Z_p \rightarrow \mathbf{N}$$

where  $\mathbf{N}$  is a set of natural numbers including zero.

Here  $Q_p/Z_p$  is a group (with respect to addition modulo 1) of numbers of the form

$$x = \sum_{i=\gamma}^{-1} x_i p^i$$

**Lemma 4.** For  $n \in Q_p/Z_p$  and  $m, k \in \mathbf{Z}$  the map  $\rho$  satisfies the conditions

$$\rho : p^m n + p^k Z_p \rightarrow p^{-m} \rho(n) + [0, p^{-k}] \quad (14)$$

$$\rho : Q_p \setminus \{p^m n + p^k Z_p\} \rightarrow \mathbf{R}_+ \setminus \{p^{-m} \rho(n) + [0, p^{-k}]\} \quad (15)$$

up to a finite number of points.

*Proof* We consider for simplicity the case  $\rho_0 = \rho$  and  $k = 0$ . Consider  $n \in Q_p/Z_p$ :

$$n = \sum_{i=\gamma}^{-1} n_i p^i$$

$$n - 1 = \sum_{i=\gamma}^{-1} n_i p^i + \sum_{i=0}^{\infty} (p-1) p^i$$



Then

$$\rho(n) = \sum_{i=\gamma}^{-1} n_i p^{-i-1}$$

$$\rho(n-1) = \rho \left( \sum_{i=\gamma}^{-1} n_i p^i + \sum_{i=0}^{\infty} (p-1) p^i \right) = \sum_{i=\gamma}^{-1} n_i p^{-i-1} + \sum_{i=0}^{\infty} (p-1) p^{-i-1} = \rho(n) + 1$$

Application of (13) for  $y = n$ ,  $y = n - 1$  proves that  $n + Z_p$  maps into  $\rho(n) + [0, 1]$ . Since the map  $Q_p/Z_p \rightarrow \mathbf{N}$  is one-to-one this proves the lemma.

**Lemma 5.** *The map  $\rho$  maps the Haar measure  $\mu$  on  $Q_p$  onto the Lebesgue measure  $l$  on  $\mathbf{R}_+$ : for measurable subspace  $X \subset Q_p$*

$$\mu(X) = l(\rho(X))$$

or in symbolic notations

$$\rho : d\mu(x) \mapsto dx$$

*Proof* Lemma 4 implies that balls in  $Q_p$  map onto closed intervals in  $\mathbf{R}_+$  with conservation of measure. The map  $\rho : Q_p \rightarrow \mathbf{R}_+$  is surjective and moreover nonintersecting balls map onto intervals that do not intersect or have intersection of the measure zero (by lemma 4). This proves the lemma.

Therefore the corresponding map

$$\begin{aligned} \rho^* : L^2(\mathbf{R}_+) &\rightarrow L^2(Q_p) \\ \rho^* f(x) &= f(\rho(x)) \end{aligned} \tag{16}$$

is an unitary operator.

Lemma 4 implies the following:

**Lemma 6.** *The map  $\rho$  maps the Haar wavelet (10) onto the function (2) (for  $p = 2$ ):*

$$\rho^* : \Psi(x) \mapsto \psi(x) \tag{17}$$

(17) is an equality in  $L^2$ : on the set of zero measure (17) may not be true.

Moreover, we have the following theorem:

**Theorem 7.** For  $p = 2$  the map  $\rho$  maps the orthonormal basis of wavelets in  $L^2(\mathbf{R}_+)$  (11) (generated from the Haar wavelet) onto the basis (5) of eigenvectors of the Vladimirov operator:

$$\rho^* : \Psi_{\gamma\rho(n)}(x) \mapsto (-1)^n \psi_{\gamma 1n}(x) \quad (18)$$

*Proof* We have

$$2^{-\gamma} \rho(x) = \rho(2^\gamma x) \quad (19)$$

Lemma 4 implies for  $\rho(n) \in \mathbf{N}$

$$\chi_{[0,1]}(\rho(2^\gamma x) - \rho(n)) = \chi_{[0,1]}(\rho(2^\gamma x - n)) \quad (20)$$

(20) is true almost everywhere. Analogously

$$\chi_{[0, \frac{1}{2}]}(\rho(2^\gamma x) - \rho(n)) = \chi_{[0, \frac{1}{2}]}(\rho(2^\gamma x - n)) \quad (21)$$

Formulas (11), (19), (20), (21) imply

$$\begin{aligned} \Psi_{\gamma\rho(n)}(\rho(x)) &= 2^{-\frac{\gamma}{2}} \Psi(2^{-\gamma} \rho(x) - \rho(n)) = \\ &= 2^{-\frac{\gamma}{2}} \Psi(\rho(2^\gamma x - n)) = 2^{-\frac{\gamma}{2}} \psi(2^\gamma x - n) \end{aligned}$$

The last equality follows from (16).

Formula (5) implies that for  $p = 2$

$$\psi_{\gamma 1n}(x) = 2^{-\frac{\gamma}{2}} \chi(2^{\gamma-1} x) \Omega(|2^\gamma x - n|_2) = (-1)^n 2^{-\frac{\gamma}{2}} \psi(2^\gamma x - n)$$

that proves (18) and finishes the proof of the theorem.

We get that after the  $p$ -adic change of variables (12) the wavelet analysis becomes the  $p$ -adic spectral analysis (expansion of a function over the eigenfunctions of the Vladimirov operator of  $p$ -adic derivation).

Using this interpretation we will call the basis (5) the wavelet basis (or  $p$ -adic wavelet basis).

Using map (16) it is possible to define the action of the Vladimirov operator in  $L^2(\mathbf{R}_+)$  by the formula

$$\partial_p^\alpha f(x) = \rho^{*-1} D^\alpha \rho^* f(x)$$

(let us note that  $D^\alpha$  and  $\rho^*$  depend on  $p$ ).

Ne can see that

$$\partial_p^\alpha f(x) = \frac{p^\alpha - 1}{1 - p^{-1-\alpha}} \int_0^\infty \frac{f(x) - f(y)}{|\rho^{-1}(x) - \rho^{-1}(y)|_p^{1+\alpha}} dy \quad (22)$$

where  $\rho^{-1}$  is the inverse map to  $\rho$ . Since  $\rho$  is not one-to-one map the map  $\rho^{-1}$  is ambiguous but ambiguity is concentrated on the set of zero measure that makes definition (22) correct.

### Acknowledgements

The author is grateful to V.S.Vladimirov, I.V.Volovich, V.A.Avetisov, A.H.Bikulov, V.P.Mikhailov and A.K.Guschin for discussion. This work was partially supported by INTAS 9900545 and RFFI 990100866 grants.

### References

- [1] V.S.Vladimirov, I.V.Volovich, Ye.I.Zelenov,  $p$ -Adic analysis and mathematical physics. World Scientific, Singapore, 1994. Russian Edition: Moscow, Nauka, 1994
- [2] V.S.Vladimirov, On the spectra of of certain pseudodifferential operators over the field of  $p$ -adic numbers, Algebra i Analiz, vol.2(1990), pp.107–124
- [3] A.N.Kochubei, Additive and multiplicative fractional differentiations over the field of  $p$ -adic numbers, in "  $p$ -Adic Functional Analysis", Lect. Notes Pure Appl. Math., vol. 192, New York: Dekker, 1997, pp.275–280
- [4] V.S.Vladimirov, Ramified characters of Idele groups of one-class quadratic fields, Proc. of the Steklov Institute of Mathematics, vol.224(1999), pp.107–114
- [5] V.S.Vladimirov, Generalized functions over the field of  $p$ -adic numbers, Usp. Mat. Nauk, vol.43(1989), No.5 pp.17–53
- [6] A.Khrennikov,  $p$ -Adic valued distributions in mathematical physics, Kluwer Academic Publ., Dordrecht, 1994

- [7] I.V.Volovich,  $p$ -Adic String, Class. Quantum Gravity, 4(1987)L83-L87
- [8] Freund P.G.O., Olson M., Nonarchimedean strings, Phys. Lett. B, 1987, Vol.199, p.186
- [9] Vladimirov V.S., Volovich I.V.,  $p$ -Adic quantum mechanics, Commun. Math. Phys., 1989, Vol.123, pp.659–676
- [10] Aref'eva I.Ya., Dragovic B., Frampton P., Volovich I.V., Wave function of the universe and  $p$ -adic gravity, Mod. Phys. Lett. A, 1991, Vol.6, pp.4341–4358
- [11] Avetisov V.A., Bikulov A.H., Kozyrev S.V., Application of  $p$ -adic analysis to models of spontaneous breaking of replica symmetry, Journal of Physics A, 1999, Vol.32, pp.8785–8791, <http://xxx.lanl.gov/abs/cond-mat/9904360>
- [12] Parisi G., Sourlas N.,  $p$ -Adic numbers and replica symmetry breaking, <http://xxx.lanl.gov/abs/cond-mat/9906095>
- [13] Carlucci D.M., De Dominicis C., On the replica Fourier transform, <http://xxx.lanl.gov/abs/cond-mat/9709200>
- [14] De Dominicis C., Carlucci D.M., Temesvari T., Replica Fourier transform on ultrametric trees and block diagonalizing of multireplica matrices, J. Phys. I France, 1997, Vol.7, pp.105-115, <http://xxx.lanl.gov/abs/cond-mat/9703132>
- [15] Daubechies I., Orthonormal bases of compactly supported wavelets, Comm. Pure Appl. Math. Vol.41 (1988) p.906; The wavelet transform, time frequency localization and signal analysis, IEEE Trans. Inform. Theory Vol.36 (1990) p.961; Ten Lectures on Wavelets, CBMS Lecture Notes Series, Philadelphia: SIAM, 1991